


Mepilex® Up

A new and innovative foam dressing designed to minimise risk of leakage¹. Inspired by patient needs and clinician perspectives.



A patented foam structure creates a dimpled surface
The innovative dimpled foam improves the capillary action, allowing fluids to spread in all directions, even against gravity.¹

The unique spreading properties minimise risk of leakage and manage exudate for several days, which may contribute to longer wear time.¹

Breathable, water-resistant backing film
Secures fluid within the dressing while allowing excess moisture to evaporate.

Safetac® wound contact layer
Our patented Safetac technology provides gentle adhesion while minimising pain and damage to the wound bed and surrounding skin²⁻⁴.

Despite its slimness, Mepilex Up can handle high volumes of fluids¹

Unique capability of spreading fluids evenly in all directions supporting better fluid handling than other foam dressings¹. This may contribute to:

Fits right into your formulary

- Highly conformable and easy to handle¹
- Able to manage exudate from high- to low-exuding wounds¹
- Longer wear time and fewer dressing changes¹
- Minimised risk of leakage¹, even under compression
- Potentially improved quality of life for patients
- Lower cost of care due to fewer dressing changes

Mepilex Up can be used on a wide range of exuding wounds, in all healing stages. It is highly adaptable and seamlessly integrated into existing protocols, without needing to change the choice of compression wraps or offloading devices.

* When comparing lab test results for Smith & Nephew Allevyn Gentle, Coloplast Biatain® Silicone NB, Convatec Aquacel® Foam, Essity Cutimed® Silitec®, according to EN13726:2023 Anne E.

Instructions for use Mepilex® Up



1. Cleanse the wound in accordance with clinical practice. Dry the surrounding skin thoroughly. Select an appropriate dressing size considering the following: For low exuding wounds, the dressing should cover the surrounding skin by at least 1 cm for small wounds and at least 3 cm for large wounds. For moderately to highly exuding wounds, regardless of wound size, the dressing should cover the surrounding skin by at least 5 cm. The dressing may be cut to suit various wound shapes and locations.



2. Remove the first release film. Apply the adherent side to the wound. Remove the remaining release film and smooth down the dressing on the skin. Do not stretch the dressing.



3. When necessary, fixate Mepilex Up with a bandage or other fixation.

Product description

Mepilex Up is a highly conformable dressing that absorbs both low and high viscous exudates, maintains a moist wound environment and minimises the risk of maceration. The soft silicone layer (Safetac®) is non-adherent to the moist wound site, but adheres gently to the dry peri-wound skin. It protects the wound and skin as well as minimises pain to patients and trauma to wounds and the surrounding skin at dressing removal.

Benefits of Mepilex Up

- The innovative dimpled foam improves the capillary action, allowing fluids to spread in all directions, even against gravity¹
- Absorbs and spreads both high- and low-viscosity fluids
- Designed to minimise leakage¹
- Minimises the risk of maceration
- Its low profile makes it highly conformable and easy to handle¹



Mepilex Up assortment (Sterile packed)

Product number	Dressing size (cm)	Pcs/box	Pcs/trp
212015	5 x 5cm	5	40
212100	10 x 10cm	5	70
212300	15 x 15cm	5	25
212200	10 x 20cm	5	25
212400	20 x 20cm	5	25
212500	20 x 50cm	2	12

Safetac® technology. Less damage. Less pain.

Safetac® adheres gently to intact skin but not the wound itself, thereby preventing damage to the peri-wound skin (e.g. skin stripping) and the wound bed, as well as minimizing associated pain on removal⁵⁻¹⁰. Safetac seals the wound edges, preventing exudate from leaking onto the surrounding skin, thus minimising the risk of maceration^{5,7,8,10,11,12}.

Indications for use

Manage exudate corresponding to low to highly exuding wounds with and without compression therapy.

Precaution(s)

- Do not use Mepilex Up on patient and/or user with known hypersensitivity to the materials/components of the product.
- Do not use together with oxidizing agents such as hypochlorite solutions or hydrogen peroxide. However, Granudacyn® has been tested and evaluated safe for use with Mepilex Up.
- If you see signs of clinical infection e.g. fever or if the wound or surrounding skin becomes red, warm or swollen, consult a health care professional for appropriate treatment.
- Do not reuse. If reused performance of the product may deteriorate, cross contamination may occur.
- Sterile. Do not use if sterile barrier is damaged or opened prior to use. Do not re-sterilise.



Scan the QR-code for further product information or visit molnlycke.xx/xxx

1. Mölnlycke Health Care. Data on File. 2024. 2. Zillmer R, et al. Biophysical effects of repetitive removal of adhesive dressings on peri-ulcer skin. J Wound Care 2006;15(5):187-191. 3. Waring M, et al. An evaluation of the skin stripping of wound dressing adhesives. J Wound Care 2011;20(9):412-422. 4. White R. A multinational survey of the assessment of pain when removing dressings. Wounds UK. 4(1). 2008. 5. Strohal R, Mittlböck M, Müller W, Hämmerle G. Management of hard-to-heal leg ulcers with an acidoxidising solution versus standard of care: The MACAN study. Journal of Wound Care. 2021;30(9):694-704. 6. Mölnlycke Health Care. Data on File. 7. White, R. A multinational survey of the assessment of pain when removing dressings. Wounds UK 2008;4(1):14-22. 8. Woo, K., Coutts, P.M., Price, P., Harding, K., Sibbald, R.G. A randomised crossover investigation of pain at dressing change comparing two foam dressings Advances in Skin and Wound Care 2009;22(7):304-310. 9. Rippon, M., Davies, P., White, R., Taking the trauma out of wound care: the importance of undisturbed healing. Journal of Wound Care 2012; 21 (8): 359-368. 10. Upton D, Solowiej K. The impact of atraumatic vs conventional dressings on pain and stress. Journal of wound care. 2012;21(5):209-15. 11. Van Overschelde, P. et al. A randomised controlled trial comparing two wound dressings used after elective hip and knee arthroplasty. Poster presentation at 5th Congress of the WUWHS, Florence, Italy, 2016. 12. Bredow J, et al. Randomised clinical trial to evaluate performance of flexible self-adherent absorbent dressing coated with silicone layer after hip, knee or spinal surgery in comparison to standard wound dressing. Poster presentation at WUWHS Congress, Italy, 2016.

Find out more at molnlycke.com

Mölnlycke Health Care AB, Box 13080, SE-402 52 Göteborg, Sweden. Phone + 46 31 722 30 00. Mölnlycke, Mepilex and Safetac trademarks, names and respective logos are registered globally to one or more of Mölnlycke Health Care Group of companies. © 2024 Mölnlycke Health Care. All rights reserved. HQIM005453

